

**TIERA**

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF IDAHO

FUSILADE® DX Herbicide

EPA REG. NO. 100-1070
EPA SLN NO. ID-040003

**FINE FESCUE GRASSES (CHEWINGS, HARD, AND CREEPING RED VARIETIES)
GROWN FOR SEED USE ONLY.**

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR PEST CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, and precautions including statements pertaining to the Worker Protection Standards, on the EPA-registered Fusilade DX Herbicide label.

This label must be in the possession of the user at the time of application.

Fine Fescues Grown for Seed

For control of downy brome (cheatgrass) (*Bromus tectorum*), quackgrass (*Agropyron repens*), bentgrass (*Agrostis sp.*), and volunteer cereals in Fine Fescue Grasses (such as Creeping Red, Chewings and Hard Fescue varieties) grown for seed in Idaho. **NOT FOR USE ON TALL FESCUE.** Refer to the federal label for a complete listing of grass weeds controlled.

Fall Application - Apply FUSILADE DX at 8 oz. per acre to actively growing downy brome in the 2 to 4 inch stage. Apply at 12 to 16 oz. per acre to actively growing quackgrass or bentgrass in the 6 to 10 inch stage. If regrowth of quackgrass or bentgrass occurs, make a second application at 12 to 16 oz. per acre before regrowth exceeds 10 inches. **Spring Application** - Apply FUSILADE DX at 8 to 16 oz. per acre to actively growing grasses. Downy brome in the 2 to 4 inch stage can be controlled by the lower rate but larger, established annual grasses (4 to 6 inches) may need the higher rate for adequate control. Apply at 12 to 16 oz. per acre to actively growing quackgrass or bentgrass in the 6 to 10 inch stage. If regrowth of quackgrass or bentgrass occurs, make a second application at 12 to 16 oz. per acre before regrowth exceeds 10 inches.

RESTRICTIONS

- Do not apply FUSILADE DX after fescue grass seedhead develops into the boot stage.
- Do not apply this product through any type of irrigation system.
- Treated fields may not be grazed or harvested for forage until after harvesting grass seed crop.
- For use only on fields of fine fescue (chewings, creeping red, and hard fescues) grown for seed. Other types of fescue may not be tolerant.
- Do not apply a total of more than 64 fl.oz of Fusilade DX per acre per year to fine fescue grasses grown for seed.

24(c) registrant:
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, NC 27419-8300

EPA SLN No.: ID-040003

Label Code: ID10700??AA0204

IDFusiladeDXFescuesSLNLbl 2-13-04